

Note: As Doppler ultrasound is the subject of the book, all entries refer to Doppler ultrasound unless otherwise indicated. Page numbers followed by “f” and “t” refer to figures and tables respectively.

A

- Abdomen (fetal), 160f
 - wall malformations, 114–115
- Abdominal diameter (AD)
 - antiphospholipid-antibody syndrome, 172f
 - IUGR assessment, 92–93, 92f
- A blood flow, 163, 163f
- A/B ratio, 26–27, 39
 - aorta (fetal), 38f
 - cerebral artery (middle), 38f, 100f
 - percentiles, 37f
 - umbilical artery, 100f
- Absent or reversed end-diastolic flow (ARED), preeclampsia, 106
- Absorption, 6
- Acardius, 122f, 126
- Acetylsalicylic acid (ASA), 47
 - preeclampsia, 105
- Acidosis, 152
 - clinical significance of Doppler ultrasound, 141
 - Doppler ultrasound, vs. cardiotocogram, 139–140, 139t
 - labor, 148
- Acoustic output, 21
- Acoustic power, 21
- Acranius, 126
- Adnexal tumor, 85, 89f
- Air, resistance to sound waves, 13
- Alias phenomenon, 18, 18f, 19f
- Allergic reactions, contrast medium, 197
- Amniocentesis, 61
- Amniotic fluid, 61
 - anhydramnios, 119, 119f, 120f
 - hydramnios-oligohydramnios, 127–128
 - pregnancy, late stages, 143
 - reduction, 56
- Amniotic fluid index (AFI), 178
- A-mode (amplitude mode), 7, 7f
- Amplitude-coded flow display, 19–20
- Anastomatoses, portal vein, 159
- Anemia
 - fetal, 61, 129–131, 130t
 - hemodynamic consequences, 129t
 - indications, 129t
 - intraperitoneal transfusion, 130–131
 - monitoring treatment, 130
 - noninvasive procedures, 129–131
 - maternal, 58
- Aneurysm(s)
 - differential diagnosis, 111
 - intra-abdominal, 115
- Anhydramnios, 119, 119f, 120f
- Annular phased array transducer, 13
- Antibody titer, 129
- Antihypertensives, uteroplacental hemodynamics, 63–64
- Antiphospholipid-antibody syndrome, 172–176, 173–176f
 - anatomical measurements, 172f
 - aorta, 175f
 - deep vein thrombosis (DVT), 172
 - ductus venosus, 176f
 - inferior vena cava, 176f
 - middle cerebral artery, 173f, 175f
 - placental-cerebral ratio, 173f
 - pulsatility index, 174f, 175f
 - resistance index, 172
 - time average maximum velocity (TAMX), 174f, 175f
 - umbilical artery, 174f
 - uterine arteries, 173f, 174f
- Aorta (fetal)
 - A/B ratio, 38f
 - antiphospholipid-antibody syndrome, 175f
 - arrhythmia, 60f
 - bifurcation, 120f
 - blood flow, 59–60, 59f
 - classification, 32
 - end-diastolic block, 48–49
 - evaluation criteria, 59
 - late pregnancy, 143–144t, 144f, 144t
 - physiological changes, 60
 - qualitative analysis, 144, 144t
 - quantitative analysis, 143, 144t
 - reference values, 59–60, 60t
 - waveform, notches, 57
 - blood redistribution, (brain-sparing), 107–108
 - correct display, 69f
 - diagnostic significance of ultrasound, 152
 - Doppler angle, 70f
 - endarteritis obliterans, 168f
 - fetal anemia, 130, 130f
 - fetal growth restriction, 167f, 177f
 - feto-fetal transfusion syndrome
 - acceptor, 180f
 - donor, 179f
 - index quotient, 37, 38f
 - multiple pregnancies, 124
 - Potter syndrome, 169f
 - pulsatility index (PI), 38f
 - renal arteries, absence, 120f
 - resistance index (RI), 38f, 49, 59t
 - near term, 148
 - second/third trimesters, 48–49, 49f
 - advantages, 49
 - bifurcation, 49f, 50f
 - diastolic reverse flow, 50f
 - mean values and standard deviation, 50f
 - resistance index (RI), 49
 - reverse flow in inferior vena cava, 50f
 - term effect, 146, 146f, 148
 - ultrasound errors, 74–75f
 - Doppler angle, 70f
- Aorta (ovarian), 41
- Aortic arch, 44f, 49f, 176f
- Aortic isthmus, 146, 147f
- Arcuate artery, ectopic pregnancy, 87–88
- Arterial pressure, fetal anemia, 131
- Arteries *see individual arteries*
- Arteriovenous shunts, 57–58
- Artifacts, 13–15
 - distal acoustic shadowing, 13, 13f
 - dorsal sound amplification, 13, 13f
 - geometrical distortion, 15
 - margin shadow, 14, 14f
 - repetition artifact, 14–15, 14f
 - side lobe, 14, 14f
 - slice thickness artifact, 14, 14f
- Ascites, 118f
- Asphyxia (fetal), 139, 148
 - cesarean section, 141
 - clinical significance of Doppler ultrasound, 141

- Asphyxia (fetal)
 Doppler ultrasound vs. cardiotocogram, 139, 139 t
- Aspirin, preeclampsia, 105
- Atenolol, 55, 64
- Atrium, 160–161, 161 f
- B**
- β_1 -sympathomimetic drugs, 55
- Balance index (BI), term effect, 146
- Balloon catheter, 198
- Baseline shift, 18, 19 f
 errors, 71 f, 73 f, 74 f, 76 f
 settings, 22
- β -blockers, uteroplacental hemodynamics, 64
- Bimodal curve, 32
- Biomechanical effects, 20–21
- Biparietal diameter (BPD)
 antiphospholipid-antibody syndrome, 172 f
 IUGR assessment, 92, 92 f
- “Birds wing” pattern, 170 f
- Bladder, 118 f, 120 f
- Blood flow
 analysis through pregnancy, 53–64
 late stages, 143–149
 breast tumors *see* Breast tumors, blood flow
 classification, 32–33, 33 f
 diastolic *see* Diastolic blood flow
 fetal anemia, 130
 fetoplacental vessels, 56–59
 indices, 27 t
 multiple pregnancies, 125
 physiological changes
 aorta (fetal), 60
 carotid arteries, 61
 fetal blood vessels, 108 f
 fetoplacental blood supply, 54, 58–59
 maternal blood vessels, 108 f
 umbilical vein, 58–59
 systolic *see* Systolic blood flow
 “term effect,” 57, 146, 146 f
 umbilical cord complications, 133, 134 f
 uteroplacental vessels, 42
 velocity measurement, 15–16, 53, 143
 indices, 26–28, 27 f, 28 f
 peak flow velocity, 161–162
- Blood pressure, uteroplacental vessels, 42
- Blood redistribution
 brain-sparing *see* Brain-sparing phenomenon
 fetal emergency, 107, 108
- Blood vessels
 adaptation in pregnancy, 42
 examination of specific vessels, 44–52
- fetal *see* Fetal blood vessels
 maternal *see* Maternal blood vessels
see also specific arteries/veins
- Blood viscosity, 25
- Blood volume
 changes through pregnancy, 145 f
 fetal anemia, 131
- B-mode (brightness mode), 8, 8 f
 record of results, 65
 safety aspects, 21
 settings, 22
 vaginal sonography, 185–186
- Bone, ultrasound energy absorption, 20
- Bradycardia, 165
- Brain-sparing phenomenon, 37, 51, 51 f, 107–108
 endarteritis obliterans, 168 f
 fetal growth restriction, 167 f
 fetal hydrocephalus, 112
 Potter syndrome, 169 f
 preeclampsia, 107–108
- Breast(s)
 circulatory changes in menstrual cycle, 202
 pulsatility index (PI), 205
 resistance index (RI), 205
 tumors *see* Breast tumors
- Breast tumors
 blood flow, 201–206
 angio mode, 203–204, 206
 benign changes, 205, 206 f
 capillaries, 201, 205, 206
 color-coded PW Doppler, 202–203, 203, 203 f
 continuous wave Doppler, 201–202, 202 f
 contrast medium, 204, 204 f
 differential diagnosis, 205–206
 malignant changes, 205, 206 f
 markers examined, 205
 power Doppler, 203–204
 pulsed wave Doppler, 202, 202 f
 blood vessels, 185 f
 malignant changes, 201, 203
 vascular supply, benign vs. malignant, 205, 206 f
- Bupivacaine, 64
- C**
- Calcium antagonists, uteroplacental hemodynamics, 64
- Cancer *see* Tumor(s)
- Capillaries, 201, 205
- Cardiac arrhythmias, 165, 165 f
- Cardiac malformations, 114, 114 f, 151
 Doppler ultrasound indication, 96
 hepatic veins and, 164 f
- Cardiotocogram (CTG), 148
- abnormal, 102
 cesarean section incidence, 105
 fetal anemia, 129
 fetal growth restriction, 177 f
 “birds wing pattern,” 170 f
 Doppler ultrasound, vs., 139–141, 139 t
 information time lead, 140, 140 f
 neonatal acidosis, 139–140, 139 t
 neonatal asphyxia, 139, 139 t
 labor, 148
 validation of indices, 100
- Carotid arteries, 51, 60–61
 bifurcation, 52 f
 internal, 61
 jugular vein, and, 52 f
 late pregnancy, 144
 low end-diastole, 52 f
 mean values and standard deviation, 52 f
 physiological flow changes, 61
 reference values, 61
- Cavitation, 20
- Center of gravity (CG), 29
- Center of gravity line (CL), 29, 30 f
- Cerebellum, intrauterine growth retardation (IURG), 93, 93 f
- Cerebral arteries
 blood flow analysis, 60–63
 diagnostic significance of ultrasound, 153
 late pregnancy, 144, 145 f
 redistribution of blood, 153
see also Brain-sparing phenomenon
 reversed diastolic flow, 113, 113 f
see also individual arteries
- Cerebral cysts, 111, 112 f
- Cerebral edema, transducer inflicted, 108
- Cerebral palsy, 151
- Cesarean section
 asphyxia, 141
 incidence, cardiotocogram (CTG), 105
- Chorion
 assessment, 88 f
 structure characteristics, 88 f
- Chorionic vessels, 42
- Chromolaparoscopy, ultrasound vs., 198
- Chromoperturbation, 198
- Circulation *see* Fetal circulation
- Clinical effectiveness, 151
- Clinical efficacy, 151
- Coccygeal teratomata, diagnosis, 118, 118 f
- Colon cancer, metastatic, 187
- Color-coded Doppler ultrasound, 19
 breast tumors, 202–203, 203 f
 color gain, 23
 endometrial carcinoma, 194 f

- fetal anomalies, diagnosis, 111
 ovaries, 188–190, 188 f, 189 t
 pulse repetition frequency, 23
 PW Doppler, 202–203, 203 f
 safety aspects, 21
 size of color window, 23
 transvaginal *see* Transvaginal ultrasound
- Color flow mapping (CFM), 186
 Concave wave front, 10, 10 f
 Conjoined twins, 121, 121 f, 122 f
 Constant flow ratio (AA), 28
 Continuous wave (CW) Doppler, 17
 breast tumors, blood flow, 201–202, 202 f
 safety aspects, 21
 Contrast medium, 197
 breast tumors, blood flow, 204, 204 f
 contraindications, 197, 204
 side effects, 197, 204
 Contrast ultrasound, 197–198
 contraindications, 197
 pain, 198
 side effects, 197
 Convex array scanner, 11, 11 f
 Corpus luteum, 87, 88, 188 f
 neovascularization, 184 f
 Curved array scanner, 11, 11 f
 Cyst(s)
 differential diagnosis, 111
 intracerebral, 111, 112 f
 kidney, 116, 117 f
 liver, 115, 116 f
 lung, 113
 ovaries, 117 f, 183 f
- D**
- Deep vein thrombosis (DVT), antiphospholipid-antibody syndrome, 172
 Demodulator, 16
 Depth gain compensation (DGC), 7
 D-galactose granules, 197
 Diabetes, Doppler ultrasound indication, 96
 Diaphragm anomalies, 113–114
 Diaphragmatic hernia, 113
 Diastolic blood flow
 breast tumors, 205–206
 diagnostic significance, 153
 end-diastolic block
 aorta, 48–49
 umbilical vessels, 57 f, 152–153
 preeclampsia, 106
 reversed
 aortic, 50 f
 cerebral arteries, 113, 113 f
 clinical significance, 156
 umbilical vessels, 55, 57 f, 152–153
- variability
 evaluation, 39
 multiple pregnancies, 125
 velocity indices, 53
 Diffraction, 6, 6 f
 Dihydralazine, 63
 Distortion, 15
 Documentation, 65–68, 66–68 f
 Doppler angle, 16, 16 f, 16 t, 69–70 f
 correction factors, 16, 16 t
 errors, 16, 16 t, 73 f, 74 f, 75 f, 76 f
 evaluation problems, 25–26
 optimal angle, 76 f
 Doppler formula, 16
 Doppler frequency (shift), 15, 15 t
 calculation, 16
 Doppler ultrasound, 15–20
 amplitude-coded flow display, 19–20
 documentation of, 65–68, 66–68 f
 errors *see* Ultrasound errors
 evaluation *see* Evaluation indices
 gynecological *see* Gynecological Doppler ultrasound
 history, 3
 instrument settings, 22–23
 during labor, 148–149
 near term findings, 143–149
 obstetric uses *see* Obstetric Doppler ultrasound
 orientation of tracings spectra, 22–23
 principles, 15–16, 15 f
 scaling time axis, 22
 wall filter *see* Wall filter
see also individual techniques; specific applications
 Doppler window, 17
 Downs syndrome, 93
 Ductus arteriosus, 62, 159, 159 f
 fetal anemia, 130 t
 Ductus venosus, 42, 159, 159 f, 161–162
 anatomy, 160, 160 f
 antiphospholipid-antibody syndrome, 176 f
 blood flow, 62–63
 abnormal, 163 f
 A blood flow, 163 f
 curve, 161 f, 162 f
 ES reduced, 164 f
 patterns, 162 f
 fetal growth restriction, 177 f
 feto-fetal transfusion syndrome
 acceptor, 180 f, 182 f
 donor, 179 f, 182 f
 measurement point, 161 f
 placental insufficiency, 164 f
- E**
- Echocardiography
 hydramnios-oligohydramnios, 128
 indications, 114
 multiple pregnancies, 125
 Eclampsia *see* Preeclampsia/eclampsia
 Ectopic pregnancy, 85–89
 characteristic findings, 85
 chorion vascularization, 183
 diagnostic validity, 87 f, 87 t
 differential diagnosis, 85–86
 early diagnosis, 184, 184 f
 perfusion resistance, 88 f
 physiological events, 86
 serum β -hCG level, 85
 transvaginal sonography, 85
 color Doppler, 86–89
 diagnostic validity, 86, 87 f, 87 t
 effectiveness, 86–87
 errors, 87–88
 Endarteritis obliterans, fetal growth restriction, 168, 168 f
 Endometrial carcinoma, 191, 194 f
 B-mode, 194 f
 color Doppler, 194 f
 detection, 193
 pulsatility index (PI), 194
 resistance index (RI), 194
 Endometrium
 atrophies, 193 f
 blood supply, 192 f
 echoic, 88 f
 examination, 191–195
 hyperplasia, 191
 luteal phase, 193, 193 f
 ovulation, 193 f
 polyps, 191, 191 f
 postmenopausal changes, 193
 premenopausal changes, 192–193
 structure, 191
 tamoxifen, 194–195
 thickness, 185, 193 f, 195
 clinical implications, 195
 tumors, 193–194
see also Endometrial carcinoma zones, 192
 Endothelium, gestational hypertension, 103
 Epidural anesthesia, 64
 Errors *see* Ultrasound errors
 Erythrocytes, velocities, 17 f
 Estrogen, endometrium perfusion, 193 f
 Evaluation indices, 25–29, 26 f, 29 f
 angle problems, 25–26
 clinical procedure, 33–34
 computerized analysis, 39
 index quotients, 37
 obstetric practice, 39

- Evaluation indices
 optical classification, 32–33
 perinatal risk and, 33 f
 pulsatile blood flow, 27 f
 qualitative measurements, 25
 quantitative measurements, 25
 reference curves, 34
 two-dimensional, 26–32
 4-point model (4 MPM), 32, 32 f
 acceleration indices, 28–29
 path length index, 29
 relative flow index, 30 f, 31–32
 temporal indices, 29
 velocity indices, 26–28, 27 f, 28 f
 waveform assessment, 30–31, 31 f
 wall filter, 26
see also specific indices
- Exercise, maternal, 60
- F**
- Fast Fourier transform (FFT), 17, 17 f
- Femoral arteries, late pregnancy sonography, 145
- Femur measurement
 antiphospholipid-antibody syndrome, 172 f
 intrauterine growth retardation (IUGR), 93, 93 f
- Fetal anemia *see* Anemia, fetal
- Fetal anomalies, 91, 111–122, 111 t
 abdominal wall, 114–115
 anhydramnios and, 119, 119 f
 Doppler ultrasound indication, 96
 head and neck region, 111–113, 111 f, 112 f, 113 f
 hydrops fetalis, 119
 malformations in early pregnancy, 121, 122 f
 placental examination, 119
see also individual anomalies
- Fetal aorta *see* Aorta (fetal)
- Fetal blood supply, 42, 43 f, 44
 fetoplacental *see* Fetoplacental blood supply
 peripheral resistance, 44
see also specific arteries
- Fetal blood vessels
 blood flow analysis, 59–63
 physiological/pathogenic profiles, 108 f
 central, 50–51
 examination, IUGR, 94
 peripheral, 47–49
 ultrasound errors, 69–82
 correct display, 69
 examination, 71–72 f
 role of Doppler angle, 69–70 f
see also Fetal blood supply; Fetal circulation; *specific arteries/veins*
- Fetal brain
 aneurysms, 112 f
 blood supply *see* Cerebral arteries
 cysts, 111, 112 f
 intracerebral pressure, 112, 112 f
 sparing *see* Brain-sparing phenomenon
- Fetal circulation, 43 f
 arterial *see* Fetal blood supply
 multiple pregnancies, 126
 near term, 143
 balance, 146–147
 venous *see* Fetal venous circulation
- Fetal disease, Doppler ultrasound indication, 96
- Fetal growth restriction, 94, 95 f, 167, 167 f
 endarteritis obliterans and, 168, 168 f
 marked, 177, 177 f
 multiple pregnancies, 124–125, 125 t
see also Intrauterine growth retardation (IURG)
- Fetal heart rate, ultrasound indications, 97
- Fetal hydrocephalus, 112–113, 112 f, 113 f
- Fetal venous circulation, 159–165
 anatomy, 159–160
 A blood flow, 163, 163 f
 clinical studies, results, 162
 display, 161–162, 161 f, 162 f
 ductus venosus to foramen ovale, 160, 161 f
 elevated intraatrial pressure and, 163
 indices, 162
 inferior vena cava to right atrium, 160–161, 161 f
 physiology, 160–161, 161 f
 premature fetuses, 164–165
see also individual veins
- Feto-fetal transfusion syndrome (FFTS), 58, 123, 127, 163, 178–181
 abnormal blood flow, 164 f
 acceptor monitoring
 aorta, 180 f
 biological measurements, 181 f
 donor vs., 182 f
 ductus venosus, 180 f
 umbilical artery, 180 f
 diagnosis, 119, 121 f
 donor monitoring
 acceptor vs., 182 f
 aorta, 179 f
 biometry, 181 f
 ductus venosus, 179 f
 umbilical artery, 178 f
- Fetoplacental blood supply, 42
 dichorionic pregnancy, analysis, 126
 examination, IUGR, 94
 late pregnancy changes, 145
 monochorionic pregnancy, analysis, 126
 multiple pregnancies, 125
see also Umbilical artery
- Fetoplacental blood vessels
 blood flow analysis, 55–59
 evaluation in preeclampsia, 106
 redistribution of blood (brain sparing), 107–108
see also Fetoplacental blood supply; Umbilical artery; Umbilical vein
- Fetoplacental perfusion disorders, screening, 102
- Fetus
 blood redistribution (brain sparing), 107–108
 examination in 3rd trimester, 91
 measurements, IUGR screening, 102
 preeclampsia effects, 106
 premature, Doppler monitoring, 164–165
see also entries beginning feto-/fetal
- Fibromas/fibroids *see* Uterine myomas
- Flow velocity waveform $F_{\max}(t)$, 25
- Focusing, 9–10
 B-mode settings, 22
 dynamic, 10
 internal, 10
- Foramen ovale, 44, 160, 160 f
- Frank–Starling principle, 60
- Frequency (f), 4
- G**
- Gain, B-mode settings, 22
- Gastrointestinal tract, malformations, 114–115
- Gastroschisis, 114, 115 f
- Geometrical distortion, 15
- Gestational age, determination, 93
- Gonadotropic-releasing hormone (GnRH), 184
- Gosling–King index *see* Pulsatility index (PI)
- Growth restriction *see* Fetal growth restriction
- Gynecological Doppler ultrasound, 183–195
 pelvic diagnosis, 185–186
 tumor angiogenesis, 184–185
see also individual techniques; specific conditions
- H**
- Head
 circumference (HC), 92, 92 f
 fetal anomalies, 111–113, 111 f, 112 f, 113 f

- Heart
 conjoined, 121 f
 disease, 98
 ectopic, 114, 114 f
 rate, 97
 recorded abnormalities, 96
 septal defects, 114, 114 f
 sparing, 108
 premature fetus, 164
see also entries beginning cardio-/cardiac
- Heat stress, blood flow changes, 54
- Height width index (HWI), 29, 29 f
- HELLP syndrome, diagnosis, 103–109
- Hemangiomas, 113, 115
- Hemodynamics *see* Blood flow
- Hepatic veins, 159–160, 159 f, 160
 A blood flow, 164 f
 cardiac malformation, 164 f
 fetal growth restriction, 177 f
 hydrops fetalis, 164 f
 negative ES blood flow, 164 f
 reverse flow, 163
- Hereditary galactosemia, 204
- Hertz (Hz), 4
- High pass filter (HPF), 19
 velocity profile loss, 31
- Hydralazine, 64
- Hydramnios-oligohydramnios, 127–128
- Hydrocephalus, 112–113, 112 f, 113 f
- Hydronephrosis, 116, 117 f
- Hydroperturbation, tubal, 184
- Hydrops fetalis, 58, 119
 hepatic veins, 164 f
 pulsations in umbilical cord, 163
- Hypertensive disorders, pregnancy, 103
see also Preeclampsia/eclampsia
- Hypervolemia, 165
- Hypoxemia, 153
- Hypoxia, umbilical vein, 58
- Hysterectomy, 191
- Hysteroscopic examination, ultrasound vs., 198
- I**
- Iliac arteries, 41, 41 f, 44, 186
- Iliac vein, 44, 45 f
- Imaging errors *see* Artifacts
- Immune system, preeclampsia, 104
- Impedance index (IMI), 28
- Inferior vena cava (IVC), 62 f, 160–161, 161 f
 antiphospholipid-antibody syndrome, 176 f
 blood flow analysis, 62
 blood flow curve, 161 f
 endarteritis obliterans, 168 f
 measurement point, 161 f
 peak blood flow velocity, 161–162
 reverse flow, 163
- Infertility, diagnosis, 197–199
- Intensity, 21
- Intensity special peak time average (ISPTA), 21
- Interference, 6
- Intervillous space, venous drainage, 42
- Intestinal hyperperistalsis, 115 f
- Intestinal peristalsis, 114
- Intracranial pressure, measurement, 112–113, 112 f
- Intrapartum pain relief, 64
- Intrauterine death, Doppler ultrasound indication, 96
- Intrauterine growth retardation (IURG), 55, 152
 biological measurements, 92–94
 basic principles, 92
 growth restriction, 94
 procedure for abnormal measurements, 93
 specific measurements, 92–93
 clinical studies, 155
 preeclampsia, 106
 screening, 102
 suspected, 94–96
- Isradipin, 64
- J**
- Jugular vein, 52 f
- K**
- Kidney
 agenesis, 117 f
 aplasia, 116
 blood supply *see* Renal artery
 polycystic, 116, 117 f
 Potter syndrome, 169 f
 pressure, 117 f
 vascular resistance, preeclampsia, 104
- L**
- Labetalol, 64
- Labor
 Doppler ultrasound during, 148–149
 pain relief, 64
 umbilical vein, effect on, 58
- Laparoscopy, 86, 88 f
- Leg, 93 f
- Linear array scanner, 11, 11 f
- Liver
 cysts, 115, 116 f
 tumor(s), 115, 116 f
 tumors, 115, 116 f
 venous drainage *see* Hepatic veins
- Low-dose Aspirin Study in Pregnancy (CLASP study), 157
- Lung anomalies, 113–114
 cystic areas, 113
 hypoplasia, 113
- M**
- Magnetostriction, 4
- Mammary tumors *see* Breast tumors
- Maternal blood vessels
 errors in Doppler ultrasound, 71–72 f
 examination methods, 44, 46–47
 IUGR, 94
 physiological and pathogenic flow profiles, 108 f
 preeclampsia, 107 f
see also specific arteries/veins
- Maternal exertion, blood flow changes, 54, 60
- Maternal heat stress, blood flow changes, 54
- Mean flow velocity $F_{max}(TA)$, 39
- Mechanical effects of ultrasound, 20–21
- Mechanical sector scanners, 12–13
 disadvantages, 13
- Menstrual cycle
 ovarian perfusion, 188
 time course of changes, 183
- Methyldopa, 63
- Metoprolol, 64
- Microstreaming, 20
- Middle cerebral artery (MCA), 50–51, 51 f, 60–61, 61 f, 111 f
 abnormal findings, 50–51, 51 f
 A/B ratio, 38 f, 100 f
 antiphospholipid-antibody syndrome, 173 f, 175 f
 cerebral edema, 61 f
 correct display, 69 f
 Doppler angle, acute, 79 f
 Doppler ultrasound, errors, 76 f
 fetal anemia, 130, 130 f
 high diastolic flow, 51 f
 index quotient, 37, 38 f
 late pregnancy, 144
 low end-diastole, 51 f
 optimal settings, 79 f
 physiological flow changes, 61
 pulsatility index (PI), 38 f, 100, 100 f
 raised intracranial pressure, 113 f
 redistribution of blood (brain sparing), 107–108
 diagnostic significance, 153
 reference values, 61
 resistance index (RI), 38 f, 100 f

Middle cerebral artery (MCA)
 reversed flow, 113 f
 validation of indices, 100

Mitral valve, fetal growth restriction, 177 f

M-mode (motion mode), 8, 9 f
 safety aspects, 21

Multiple pregnancies, 123–128
 acardius acranium, 121, 122 f, 126
 circulatory communication disorders, 127
 clinical significance of Doppler ultrasound, 156
 conjoined twins, 121, 121 f, 122 f
 cord looped round neck, 126
 diagnostic procedures, 128 f
 dichorionic, 123, 123 f
 Doppler ultrasound indications, 96
 growth restrictions, 124–125, 125 f
 fetofetal transfusion syndrome, 127
 hydramnios-oligohydramnios, 127–128
 monoamniotic, 123, 123 f
 monochorionic, 127
 mortality, 156 t
 resistance index (RI), 124
 S/D ratio, 124
 special considerations, 126–128
 studies, 124–126
 theoretical considerations, 125–126
 systolic/diastolic variability, 125
 twin reversed arterial perfusion (TRAP), 123, 126
 twin-to-twin transfusion syndrome
see Fetofetal transfusion syndrome (FFTS)

umbilical vessels, 48, 56
 vasa previa, 126
 velamentous insertion, 126

N

Neck, fetal anomalies, 111–113

Neoplasia *see* Tumor(s)

Nitrogen monoxide (NO), flow changes, 55

Noise, errors in Doppler ultrasound, 73 f, 75 f

Nordic trial, 193

Notching, importance, 46, 46 f

Nutritional impairment, multiple pregnancies, indications, 124–125

Nyquist limit, 18, 19, 19 f
 settings, 22

O

Obstetric Doppler ultrasound
 cases illustrating use, 167–182
 clinical significance of Doppler ultrasound, 151–156, 154 t
 analysis of Doppler studies, 154–155
 conclusions, 156–157
 cumulative metaanalysis, 155–156
 diastolic reverse flow, 156
 diagnostic significance, conclusions, 156–157
 diagnostic significance of, 151–153
 evaluation indices, 39
 indications for Doppler ultrasound, 91–98
 eclampsia, 96
 intrauterine death, 96
 IURG and biological measurements, 92–94
 malformations suspected, 91
 preeclampsia, 96
 status post dysmature, 96
 screening at-risk populations *see* Obstetric screening of at-risk groups
see also Pregnancy; *individual techniques; specific conditions*

Obstetric screening of at-risk groups, 99–102
 cost-benefit ratio, 100
 fetoplacental perfusion disorders, 102, 102 f
 intrauterine growth retardation (IURG), 102, 102 f
 populations, 101
 preeclampsia, 101–102
 preventative care and, 171, 171 f
 uteroplacental perfusion disorders, 101–102
 validity of indices, 100–101, 100 f, 101 f
 validity of tests, 99–100

Occipitofrontal diameter (OFD), 92, 92 f

Omphalocele, 114, 115 f

Optical classification, 32–33

Oscillation, 3–4

Ovarian artery, 186

Ovarian carcinoma, 187, 190 f
 angiogenesis, 188
 color-coded Doppler ultrasound, 188–190
 preoperative staging, 189 t
 Doppler index values, 189 t
 preoperative staging, 187 t
 Schillinger score, 187 t

Ovaries, 186
 color-coded Doppler ultrasound, 188–190, 188 f, 189 t

cystic, 117 f, 183 f
 diagnostic procedures, 187–190, 187 f
 ectopic pregnancy, 183 f
 evaluative criteria, 187
 size, 187
 structure, 187
 menstrual cycle changes, 188
 perfused, 188 f
 resistance index (RI), 188
 test validity differentiation, 189 t
 tumors *see* Ovarian carcinoma

Ovulation, endometrium, 193 f

Oxytocin challenge test (OCT), 152

P

Parallel scanner, 11, 11 f

Path length index (PLI), 29

Pelvis
 diagnostic procedures, 185–186, 186 f
 evaluation, 186
 qualitative, 186
 quantitative, 186
 indices, 186
 vascular perfusion, 183 f

Penetration depth, B-mode settings, 22

Perinatal mortality, 156 t, 157
 reverse flow and, 56–57

Phased array scanner, 11, 12 f

Piezoelectric ceramic, 5 f

Piezoelectric crystal, 5

Piezoelectric effect, 5

Placenta
 anomalies, 119
 clinical significance, 141
 arteriovenous shunts, 57–58
 biopsy, 170 f
 blood flow changes, 55
 blood supply
 fetoplacental *see* Fetoplacental blood supply
 uteroplacental *see* Uteroplacental blood supply
 vascular bed, 55, 104, 170 f

cotyledon, 42
 insufficiency, 91
 ductus venosus, 164 f
 mass, late pregnancy, 143
 Potter syndrome, 170 f
 premature separation, 57
 unilateral implantation, 152
see also Fetoplacental blood vessels; Uteroplacental blood supply

Placental-cerebral ratio (PCR), anti-phospholipid-antibody syndrome, 173 f

Polycystic kidney, 116, 117 f

Portal sinus (sinus venous), 42, 159

- Portal vein (PV), 159, 159 f
 Potter syndrome, 120 f, 169
 exclusion of, 169–170 f
 Preeclampsia/eclampsia, 152
 diagnosis, 103–109
 evidence for fetal risk, 106–106
 uteroplacental arteries, 105
 Doppler ultrasound indications, 96,
 104
 flow changes, 55
 IUGR risk, 106
 notching, 46, 46 f
 pathological changes, 103–104
 screening, 101–102
 Pregnancy
 blood flow analysis, 53–64
 cerebral arteries, 60–61
 fetal vessels, 59–63
 fetoplacental vessels, 55–59
 therapeutic measures, effects,
 63–64
 uteroplacental vessels, 53–55,
 53 f
 delivery date, 148
 early, fetal malformations, 121–122
 ectopic *see* Ectopic pregnancy
 first trimester ultrasound, 96
 multiple *see* Multiple pregnancies
 near term findings, 143–149
 circulatory balance, 146–147
 clinical conclusions, 147–148
 Doppler ultrasound evaluation of
 acute changes, 147–148
 maternal adaptation, 143
 physiological, 143–145
 second trimester ultrasound, 96
 term and postterm findings, 145–
 147, 146 f, 146 t
 third trimester ultrasound, 96
see also Obstetric Doppler ultra-
 sound; *specific complications*
 Pregnancy-induced hypertension *see*
 Preeclampsia/eclampsia
 Prostacyclin, 63
 preeclampsia, 103
 Prostaglandins, uteroplacental
 hemodynamics, 63
 Pulsatility index (PI), 27 f, 31, 39
 antiphospholipid-antibody syn-
 drome, 174 f, 175 f
 aorta (fetal), 38 f, 144
 breast, 205
 cerebral artery (middle), 38 f, 100,
 100 f
 ectopic pregnancy, 87
 endometrial carcinoma, 194
 fetal growth restriction, 167 f
 fetal hydrocephalus, 112
 intersection of medians, 38 f
 maternal position, 54
 reference curves, 35 f
 umbilical artery, 100 f
 uterine myomas, 190 t
 Pulsed wave (PW) Doppler, 17–18,
 18 f
 alias phenomenon, 18, 18 f, 19 f
 breast tumors, blood flow, 202,
 202 f
 safety aspects, 21
 settings, sample volume, 22
 Pulse-echo procedure, 7
 Pulse repetition frequency (PRF), 17–
 18
 baseline shift, 18, 19 f
 settings, 22, 23
- Q**
 Quartz, 5
- R**
 Radial artery, 41
 Record keeping, 65–68, 66–68 f
 Reference curves, 34, 35 f
 Reference values, 60 t
 aorta (fetal), 59–60, 60 t
 carotid arteries, 61
 middle cerebral artery (MCA), 61
 renal arteries, 62
 umbilical blood flow, 55, 58, 60 t
 uteroplacental blood flow, 54, 60 t
 Reflection, 5–6, 5 f
 Refraction, 5–6
 Region of interest (ROI), 19, 19 f
 Relative acceleration time (RAT), 29
 Relative flow index (RFI), 30 f, 31–32
 Relative mean inflow time (rMIT), 29
 Relative (receiver) operating charac-
 teristic (ROC) curve, 99, 99 f
 test validities, 100 f, 101 f
 Relative rate flow index (RI), 30
 Relative spectral area, 31
 Renal agenesis, 117 f
 Renal aplasia, 116
 Renal artery
 bifurcation, 120 f
 blood flow analysis, 61–62, 62 f
 evaluation criteria, 62
 examination, 116, 117 f
 late pregnancy sonography, 145
 Potter syndrome, 169 f
 reference values, 62
 Renal vascular resistance, preeclam-
 psia, 104
 Repetition artifact, 14–15, 14 f
 Resistance index (RI), 26–27, 39
 antiphospholipid-antibody syn-
 drome, 172
 aorta (fetal), 38 f, 49
 late pregnancy, 144
 endometrial carcinoma, 194
 fetal intracranial flow curves, 145
 middle cerebral artery, 38 f, 100,
 100 f
 multiple pregnancies, 124
 ovaries, premenopausal, 188
 percentiles, 36 f
 term effect, 146 f
 tumors, 185
 umbilical artery, 55, 100 f
 uterine myomas, 190 t
 uteroplacental artery, 54, 54 f
 Resolution, 9
 Reverberation artifact, 14–15, 14 f
 Rotary transducer, 12, 12 f
- S**
 Safety aspects, 20–21
 Saline infusion sonography, 194
 Scanning procedures, 10–13
 curved/convex array scanner, 11,
 11 f
 linear array scanner (parallel), 11,
 11 f
 mechanical sector scanners, 12–13
 phased array scanner, 11, 12 f
 principle of operation, 10–11, 10 f
 sector scanner, 11, 11 f
 Scattering, 6, 6 f
 Sector scanner, 11, 11 f
 Serum β -hCG level, ectopic preg-
 nancy, 85
 Shadowing, 13, 13 f, 14, 14 f
 distal acoustic, 13, 13 f
 margin, 14, 14 f
 Side lobe artifact, 14, 14 f
 Skull
 blood supply, 111
 measurement
 antiphospholipid-antibody syn-
 drome, 172 f
 IUGR assessment, 92
 Slice thickness artifact, 14, 14 f
 Sound amplification, dorsal, 13, 13 f
 Sound field, 9–10
 focusing, 9–10
 resolution, 9
 Sound waves, 3–4, 3 f
 absorption, 6
 air, resistance to, 13
 concave wave front, 10, 10 f
 diffraction, 6, 6 f
 frequency (f), 4
 frequency ranges, 4 f
 generation, 4–5
 interference, 6
 reflection, 5–6, 5 f
 refraction, 5–6
 scattering, 6, 6 f
 substance density effects, 4 t
 velocity, 4
 wavelength (λ), 4

Spectrum broadening index (SBI), 28, 28 f, 39
 Spectrum broadening ratio (SBR), 28, 28 f, 39
 Spiral artery, 192
 ectopic pregnancy, 87–88
 Sufentanil, 64
 Supine position, blood vessel compression, 56
 Supraventricular extrasystoles (SVES), 165, 165 f
 Supraventricular tachycardia (SVT), 165, 165 f
 Systolic blood flow
 acceleration, measurement, 39
 breast tumors, 205–206
 variability
 evaluation, 39
 multiple pregnancies, 125
 velocity
 aorta, 59
 fetal anemia, 130
 Systolic decay time index (SDTI), 29, 29 f
 Systolic delay, measurement, 39

T

Tamoxifen, 191
 endometrial carcinoma, 194
 “Term effect,” 57, 146, 146 f
 Thermal effects of ultrasound, 20
 Time average maximum velocity (TAMX), 174 f, 175 f
 Time gain compensation (TGC), 7
 Time-weighted pulsatility index (twPI), 29
 Tourmaline, 5
 Toxemia of pregnancy *see* Preeclampsia/eclampsia
 Tracings, orientation of, 22–23
 Transducers *see* Ultrasonic transducers
 Transient cavitation, 20
 Transvaginal ultrasound, 85, 197–199
 color Doppler, 88 f
 blood flow curve, 89 f
 critical evaluation, 89
 ectopic pregnancy, 86–89, 87 f, 87 t
 Tricuspid valve, fetal growth restriction, 177 f
 Tumor(s)
 angiogenesis, 184–185
 assessment, 183
 benign, 201
 blood flow, 185
 blood vessels, 184, 185, 185 f, 201
 diagnostic criteria, 185
 growth, 185
 resistance index (RI), 185
 postsystolic notch, 57 f
 reference values, 55, 61 t
 reverse diastolic flow, 57 f
 blood redistribution (brain-sparing), 108
 cardiotocogram, 102
 clinical significance of ultrasound, 154–156
 diagnostic significance of ultrasound, 152–153
 display
 color, 47 f
 correct, 69 f
 Doppler angle, 73 f, 77 f
 endarteritis obliterans, 168 f
 fetal growth restriction, 167 f, 177 f
 feto-fetal transfusion syndrome acceptor, 180 f, 182 f
 in donor, 178 f, 182 f
 multiple pregnancies, 124, 125
 neonatal asphyxia, 139, 139 t
 normal, 47
 pH values, 141 t
 Potter syndrome, 169 f
 preeclampsia effects, 107 f
 resistance index (RI), 55, 148
 spectral Doppler, 48
 ultrasound errors, 77 f
 baseline, 73 f
 Doppler angle, 73 f
 wall filter error, 73 f
 Umbilical blood vessels, 42 f
 blood flow analysis, 55–59, 55 f
 compression, 56
 reverse flow, 56–57
 see also Umbilical artery; Umbilical vein
 Umbilical cord, 44 f
 anomalies, 119, 121 f
 blood vessels *see* Umbilical blood vessels
 complications, 133–136, 134 t
 coils under fetus, 78 f
 hemodynamics affected, 133, 134 f
 knot, 134 f
 looped round neck, 126, 133 f
 puncture, 157
 gastrointestinal malformations, 114
 gastroschisis, 115 f
 omphalocele, 115 f
 perfusion, fetal anemia, 131 f
 pulsations, hydrops fetalis, 163
 reverse flow, 78 f
 systolic notch, 134 f
 thickness, 133
 tumor, 121 f
 tumor(s), 121 f
 Umbilical vein, 42, 58 f, 159, 159 f
 blood flow analysis, 58–59
 diastolic block, 152–153
 flow patterns, 162, 162 f

U

Ultrasonic transducers, 9 f
 annular phased array, 13
 cerebral edema caused by, 108
 rotary, 12, 12 f
 selection, 22
 wobbler, 12 f, 13
 Ultrasound, 3–23
 artifacts *see* Artifacts
 definitions, 21
 diagnostic values, 4
 errors *see* Ultrasound errors
 fetal anemia, 129
 history, 3
 image generation, 7–8
 instrument settings, 22–23
 physical effects, 5–6, 5 f, 6 f
 procedures *see* Scanning procedures
 record of results, 65–68
 safety aspects, 20–21
 scanning procedures, 10–13
 see also Sound waves; Ultrasonic transducers; *specific techniques*
 Ultrasound errors, 69–82
 baseline shift, 71 f, 73 f, 74 f, 76 f
 Doppler angle, 16, 16 t, 69–70 f, 73 f, 74 f, 75 f, 76 f
 examination, 71–72 f
 imaging *see* Artifacts
 noise, 73 f
 transvaginal color Doppler, 87–88
 wall filter, 73 f, 74 f
 Umbilical artery, 47–48, 73 f
 abnormal, 47, 102
 A/B ratio, 100 f
 antiphospholipid-antibody syndrome, 174 f
 ascites, 118 f
 blood flow analysis, 55–58, 55 f
 diastolic flow, 55
 end-diastolic block, 57 f, 152
 end-diastolic shift, 32
 morphological flow changes, 56–58
 pathological flow changes, 56
 physiological flow changes, 56

- physiological/pathological changes, 58–59
 - reverse flow, 152–153
 - blood flow, volume, late pregnancy, 145 f
 - diagnostic significance of ultrasound, 152–153
 - diameter, 58
 - display, correct, 69 f
 - examination, 161
 - fetal anemia, 130
 - intraoperative transfusion monitoring, 130–131
 - pulsations, 58, 58 f, 127, 162–163, 163 f
 - diagnostic significance, 153
 - patterns, 162
 - reference values, 58
 - Urogenital system
 - anomalies, 116, 117 f, 118 f
 - tumors, 116
 - Uterine arteries, 53 f, 71–72 f, 77 f, 192
 - antiphospholipid-antibody syndrome, 173 f, 174 f
 - correct display, 69 f
 - Doppler angle, 69–70 f
 - fetal growth restriction, 167 f
 - flow profile, 71 f
 - high risk patients, monitoring, 171 f
 - normal flow, 77 f
 - notched, 53 f, 72 f
 - postsystolic notch, 53 f
 - Potter syndrome, 169 f
 - preeclampsia effects, 107 f
 - pulse repetition frequency
 - display, 77 f
 - errors, 71 f, 72 f
 - reverse flow, 71 f
 - second trimester examination, 44, 45 f, 46, 46 f
 - abnormal findings, 44, 46–47
 - high risk patients, 171 f
 - iliac to arcuate artery, 45 f
 - notching, 46 f
 - origin from iliac artery, 41 f
 - transvaginal display, 46 f
 - superimposed on iliac, 53 f
 - third trimester examination, 47
 - high risk patients, 171 f
 - ultrasound errors, 69–82
 - amplification, 72 f
 - baseline, 71 f
 - notches, 53 f, 72 f
 - poor focus, 70 f
 - pulse repetition frequency, 71 f, 72 f
 - Uterine blood flow
 - correct display, 69
 - role of Doppler angle, 69–70 f
 - ultrasound errors, 69–82
 - see also* Fetal blood supply; Umbilical artery; Uterine arteries; *specific vessels*
 - Uterine contractions, blood flow changes, 54
 - Uterine myomas, 184, 190–191, 190 f
 - Doppler indices, 190 t
 - Uterine tube, 184 f, 197–199
 - Uterine tumors, 191
 - Uterine vascular resistance, preeclampsia, 104
 - Uteroplacental arteries *see* Uteroplacental blood supply
 - Uteroplacental blood supply, 41–42, 41 f
 - clinical significance of ultrasound, 154
 - diagnostic significance of ultrasound, 152
 - endarteritis obliterans, 168 f
 - flow analysis, 53–55 53 f
 - abnormal flow changes, 55
 - indices of systolic and diastolic velocity, 53
 - medications, 55
 - physiological flow changes, 54
 - postsystolic notch, 53
 - reference values, 54, 60 t
 - impairment, multiple pregnancies, 125
 - perfusion, 41 f
 - preeclampsia, 105, 106
 - therapeutic evaluation, 63–64
 - antihypertensives, 63–64
 - epidural anesthesia, 64
 - prostaglandins, 63
 - see also individual arteries*
 - Uteroplacental perfusion
 - disorders, screening, 101–102
 - fetal growth restriction, 167 f
 - Uteroplacental unit, 41–52
 - examination methods, 44–52
 - maternal vessels, 44, 46–47
 - peripheral fetal vessels, 47–48
 - near term findings, 143
 - vascular supply, 41–44, 43 f
 - fetal blood supply, 42, 43 f, 44
 - fetoplacental *see* Fetoplacental blood supply
 - uteroplacental *see* Uteroplacental blood supply
 - Uterus
 - blood vessels, 185 f, 192 f
 - see also specific vessels*
 - endometrium *see* Endometrium
 - myomas (fibromas/fibroids) *see* Uterine myomas
 - removal, 191
 - scanty serometra, 186 f
 - see also entries beginning utero-/uterine*
- V**
- Vaginal examination, 185–186
 - Vasa previa, 126
 - Vascular lesions, flow changes, 55
 - Vascular perfusion, 205
 - Vascular proliferation, tumors, 201
 - Vascular spasm, preeclampsia, 103, 104 f
 - Velamentous insertion, 126
 - Velocity, relative reduction, 39
 - Vena cava, inferior *see* Inferior vena cava (IVC)
 - Venous pressure, fetal anemia, 131
 - Ventricular extrasystoles (VES), 165
- W**
- Wall filter, 19
 - errors and, 73 f, 74 f
 - evaluation indices, 26
 - settings, 22
 - Waveform indices, aorta, 59–60
 - Wavelength (λ), 4
 - Wobbler transducer, 12 f, 13